

## Original Article

# Effect of methanolic crude extract of *Aframomum melegueta* (A.m) seeds on selected lactogenic hormones of Albino rats

Hyacinth Ibekwe

Department of Animal Science, Cross River University of Technology (CRUTECH) Obubra Campus, Nigeria

Received January 10, 2019; Accepted April 29, 2019; Epub June 15, 2019; Published June 30, 2019

**Abstract:** The search for alternative control of excessive milk secretion associated with exclusive breast feeding necessitated this study. This study was designed to investigate the effect of methanolic crude extract of *Aframomum melegueta* (A.m) seed by measuring serum concentrations of selected lactogenic (prolactin, estradiol and progesterone) hormones in lactating and non-lactating Albino rats. A total of 18 non-lactating and lactating rats each were assigned into three (3) treatments of 6 rats each. Treatment 1 and 2 (controls) received 0.11 mg/kg/day of reference drug (bromocriptine) and 100 mg/kg/day of Dimethylsulfoxide (DmsO) 1 ml:9 ml of normal saline which is referred to as normal control in this study. Treatment 3 received 100 mg/kg/day of A.m methanolic crude extract. Serum samples were collected and analyzed for prolactin, estradiol and progesterone by ELISA techniques. The result obtained showed a 21.13% and 25.12% significant ( $P < 0.05$ ) reduction in serum concentration of prolactin for both non-lactating ( $26.15 \pm 0.15$  ng/ml) and lactating ( $14.90 \pm 1.16$  ng/ml) rats relative to ( $33.23 \pm 1.82$  and  $19.90 \pm 1.16$  ng/ml) normal control rats but a significant ( $P < 0.05$ ) increase ( $26.15 \pm 0.15$  ng/ml;  $14.90 \pm 1.16$  ng/ml) in serum levels of the same hormone relative to ( $12.56 \pm 0.89$  ng/ml;  $10.56 \pm 0.29$  ng/ml) reference control drug (bromocriptine) in both groups. Serum estradiol levels were significantly ( $P < 0.05$ ) reduced by methanol extract relative to normal control rats in both groups. There was however, no significant ( $P > 0.05$ ) changes in serum levels of estradiol relative to reference control drugs in both rats. Serum levels of progesterone were not significantly altered in both groups relative to normal control rats. The above findings confer on the extract antilactogenic capability and hence a good alternative to bromocriptine.

**Keywords:** Lactogenic hormones, methanol extract, *Aframomum melegueta*, Albino rats

## Introduction

Breast milk production is a complex physiologic process involving physical and emotional factors and the interaction of multiple hormones, the most important of which is believed to be prolactin [1]. With parturition and expulsion of placenta, progesterone level falls and full milk supply is initiated i.e. lactogenesis [2]. In most laboratory species, mammary gland lobuloalveolar growth could be induced after ovariectomy with combination of estrogen and progesterone injection [3]. Extensive proliferations of the mammary gland in response to ovarian hormones occur only if the pituitary gland is intact. In both the rat and the mouse, studies revealed the hormonal requirements for mammary gland proliferation in the absence of endogenous

mammogenic hormones i.e. after the removal of the pituitary and the ovaries or the pituitary, ovaries and adrenals [4]. Studies by [5] showed that in the hypophysectomized-ovariectomized-adrenalectomized rat, the pituitary hormone by themselves in sufficient doses were capable of inducing tubuloalveolar development [3]. This observation does not imply that the ovarian hormones are of no importance in normal mammary growth but that the pituitary hormones must be regarded the more essential since the ovarian hormones in the absence of pituitary hormones have little or no mammogenic activity in normal mammary growth [3].

The pituitary hormone of interest in this study is prolactin which causes milk secretion from the breast after estrogen and progesterone prim-

## Methanolic extract of A.m on lactogenic hormones

ing. Its effect on the breast involves increased action of messenger ribonucleic acid (mRNA), increased production of casein and lactalbumin [6]. Increased serum concentration of prolactin during pregnancy causes enlargement of the mammary gland and increase the production of milk [7]. The high level of progesterone during pregnancy acts directly on the breast to stop ejection of milk [8]. During pregnancy, high circulating concentration of estrogen promotes prolactin production [9]. The secretion is however, inhibited by hypothalamic and section of pituitary stalk which leads to an increase in circulating prolactin [10]. Prolactin levels peak at rapid eye movement sleep (REM) and in the early hours of the morning [11].

The use of synthetic drugs to improve milk secretion in exclusive breast feeding has more often been challenged with hyperprolactinemia, prolactinoma, lactation amenorrhoea and galactorrhoea [12]. Lactogogues or lactogenic agents are medication of other substances capable of initiation, maintenance or augmentation of maternal milk production [13]. The galactogogues in common use include metoclopramide, domperidone and sulpiride which act by blocking dopamine receptors and therefore, the prolactin inhibiting action of dopamine. Domperidone is said to increase milk secretion and is safe [14]. The continued use of these drugs with their attendant side effects led to the discovery and use of antilactogenic agents. The orthodox antilactogenic drugs (cabergoline, pergolide, bromocriptine, quinagolide) required to handle the emerging side effects are associated with serious side effects (nausea, fatigue, postural hypotension, nasal congestion, exacerbation of psychosis, seizure and cardiovascular injuries that limits completion of dosage regimen. It therefore becomes imperative to seek for natural alternative with little or no adverse effect on the recipients.

Aframomum melegueta is the plant of choice. Aframomum melegueta is an herbaceous perennial plant native to swampy habitats along the West African Coast. Its trumpet-shaped purple flowers develop into 5-7 cm long pods containing small reddish-brown seeds [15]. Essential oils occur only in traces [16]. The use of Aframomum melegueta root extract in inhibiting excessive milk secretion by nursing mothers has been reported [17]. The dose dependent inhibitory effect of Vincristine on milk

secretion in goat was reported with 1 mg of this alkaloid decreasing milk yield to approximately 40% of previous yield [18]. Phytochemistry revealed the presence of alkaloids, glycosides, tannin, flavonoids, sterols, triterpenes and oils [19]. In addition, seed extract of A. melegueta contains polyphenols and hydrocyanic acid.

This experiment is therefore; set to investigate the effect of the seed of Aframomum melegueta methanol crude extract in Albino rats using bromocriptine as reference control drugs. Its action on the lactogenic hormones (prolactin, estradiol and progesterone) is the interest of this research.

### Materials and methods

#### *Preparation of plant material*

Fresh fruits of Aframomum melegueta (A. melegueta) weighing 500 g were purchased from Apiapum market in Obubra Local Government Council of Cross River State of Nigeria. The fruits were sundried for two weeks to facilitate removal of the seeds from the pods (capsule). The seeds were further sundried to constant weight within four days and stored in an airtight flask for extraction.

#### *Preparation of Aframomum melegueta seed extract*

The seeds were ground to powdered form by the use of motor powered milling machine at Root Crop Research Institute, Umudike in Abia State. A 250 g weight of Aframomum melegueta seed powder was first extracted with petroleum ether (60-80°) C to remove the fat. The ether extract was carefully decanted leaving the residue for further extraction in methanol. The soxhlet apparatus was filled with methanol and 200 g of the residue was wrapped in a thimble and placed in a soxhlet extractor fitted to a 500 ml round bottom flask seated on a hot plate. The reflux condenser was attached to the extraction tube and power was supplied by switch linking the electric hot plate cable. The methanol vapour passes up the side tube of condenser and runs back on to the residue in the thimble. After siphoning over for 24 times, the experiment was stopped just before the next lot of methanol was at the point of siphoning over. The methanol soluble fraction or methanolic extract was then preserved at 4°C ready for administration to experimental rats.

## Methanolic extract of A.m on lactogenic hormones

### *Preparation of stock solution of crude extract of Aframomum melegueta seeds*

One gram (1 g) of methanol extract of Aframomum melegueta seed was dissolved in one (1 ml) milliliter of Dimethyl sulfoxide (DmsO) and suspended in 9 ml of normal saline (0.9% NaCl). The stock solution was immediately administered to rats in treatment III of this study.

### *Animal and animal procurement*

A total of 18 wistar rats each for non-lactating and lactating groups were assigned on the basis of weight (150-200 g) into 3 treatments of 6 rats each. The animals were housed in the Department of Biochemistry, University of Calabar under standard laboratory conditions of ambient temperature of 26°C and adequate ventilation with relative humidity of 50% and a 12 hour day-light cycle. Animal ethic regulations were observed as per the Institution's animal ethics regulation. The administration of Aframomum melegueta seed extract commenced 3 days postpartum and lasted 3 weeks for both non-lactating and lactating rats. Rats in treatment group I: received bromocriptine (reference standard drug) at a dose of 0.11 mg/kg/day according to manufacturer's instruction. Treatment group II rats received 100 mg/kg/day of dimethyl sulfoxide, DmsO (1 ml in 9 ml of normal saline) for normal control treatment. The rats in treatment III received 100 mg/kg/day of methanol seed extract of Aframomum melegueta via oral intubation. The treatment was terminated after a 3 week period of administration of crude extract. The rats were starved of food overnight, and sacrificed by decapitation. The blood sample was aseptically collected via cardiac puncture and transferred into sample labeled bottles, while the heart was still beating. The collected blood was allowed to stand for 2 hours to perfect clotting of blood and centrifuged at 1000 rpm and serum removed with Pasteur pipette for assay of prolactin, estradiol and progesterone concentrations.

### *Estimation of serum prolactin concentration*

This was carried out by microwell enzyme immuno-assay using the method of [20]. A total of 41 streptavidin located micro-plate was removed from the zip-lock kit being-for a substrate and two each for reference standard and

control and 36 micro-plates for the test samples. They were properly labeled and placed in a micro-plate holder. Twenty five (25 µl) micro-liter of standards, controls and samples were dispensed into each micro-well. A total of 100 µl of conjugate reagent was dispensed into each well and thoroughly mixed for 30 seconds. These were incubated at room temperature for 60 minutes and the content of the well discarded by decantation and the micro-plate blotted dry with absorbent paper. A 300 VI wash was aspirated and repeated twice making a total of three washes. One hundred (100 µl) micro liter of 3,3',5',5'-tetramethyl bendine (TMB) was added into each well and gently mixed for 110 seconds. These were incubated at room temperature in the dark for 15 minutes. A 50 µl of stop solution was gently added to each well and mixed after the blue colour of the well has completely changed to yellow. The absorbance of each well was read at 450 nm wave length. The average absorbance values were calculated from the duplicate standards, controls and sample and a standard curve was constructed by plotting the average absorbance obtained from each reference standard against its assigned concentration in ng/ml on linear graph paper with the absorbance on the vertical (Y) axis and the concentrations on the horizontal (x) axis. The corresponding concentration of prolactin in the sample was determined from the intersecting point of the curve.

### *Estimation of serum estrogen (estradiol) concentration*

This was determined by microwell enzyme immunoassay according to the method of [21], and progesterone serum concentration according to the method of [22].

### *Data analysis*

Data collected were subjected to statistical analysis using one way comparison analysis of variance (ANOVA). A one way analysis of variance was used to compare differences among groups of treatment and significant means separated by least significant difference. The least significant difference (LSD) indicates how different the difference is among the means. This was statistically computed using this formular,  $LSD = t_{\alpha, 0.25} \text{ (error degree of freedom)} \times \frac{\sqrt{2S_e^2}}{r}$ . Where  $S_e$  = Error mean

## Methanolic extract of A.m on lactogenic hormones

**Table 1.** Treatment schedule for experimental rats

Animal category	Treatments or design of experiment		
	Bromocriptine	Control	Methanol extract
Non-lactating (NL) rats	6	6	6
Lactating (L) rats	6	6	6

square and  $r$  = Number of observation per treatment.

### Result

Serum concentrations of selected lactogenic (prolactin, estrogen and progesterone) hormones were presented in **Table 2**. The result indicates that bromocriptine (standard reference drug) produced a 62% and 52% significant ( $P < 0.05$ ) decrease in serum concentration of prolactin ( $12.56 \pm 0.89$  ng/ml) relative to ( $33.23 \pm 1.82$  ng/ml,  $26.15 \pm 0.15$  ng/ml) control and methanol extract treated non-lactating rats respectively. The standard reference drug (bromocriptine) produce a 7.87% significant but marginal ( $P > 0.05$ ) decrease in serum concentration of estradiol ( $2.81 \pm 0.12$  ng/ml) with respect to control ( $3.05 \pm 0.06$  ng/ml) and a 4.63% ( $2.68 \pm 0.08$  ng/ml) methanol extract treated non-lactating rats. The serum concentration of progesterone was however, significantly ( $P < 0.05$ ) reduced to 60.44% ( $127.96 \pm 6.06$  ng/ml) by bromocriptine in comparison with control ( $323.43 \pm 0.20$  ng/ml) and a 2.4% non-significant reduction by methanol extract treated ( $315.72 \pm 7.72$  ng/ml) non-lactating rats. Methanol extract produced a 12.13% significant ( $P < 0.05$ ) reduction in serum concentration of estradiol ( $2.68 \pm 0.08$  ng/ml) with respect to control ( $3.05 \pm 0.06$  ng/ml) but did not however, produce significant ( $P < 0.05$ ) reduction in serum concentration of progesterone ( $315.72 \pm 7.72$  ng/ml) relative to control ( $323.43 \pm 0.20$  ng/ml) nonlactating rats. For the lactating rats, in **Tables 1-3**, administration of bromocriptine brought about a 47.00% significant ( $P < 0.05$ ) lowering ( $10.56 \pm 0.26$  ng/ml) of prolactin in comparison with the control and 29.13% methanol extract ( $19.90 \pm 1.16$  ng/ml;  $14.90 \pm 1.16$  ng/ml) respectively. Methanol extract also caused a 25.13% significant ( $P < 0.05$ ) reduction in serum prolactin ( $14.90 \pm 1.16$  ng/ml) relative to control ( $19.90 \pm 1.16$  ng/ml). For serum concentration of estradiol, bromocriptine as well as methanol extract produced 24% significant ( $P < 0.05$ ) reduction ( $2.41 \pm 0.80$ ,  $2.40 \pm 0.03$  ng/ml) relative to control ( $3.17 \pm 0.09$  ng/ml) treat-

ment but there was no significant ( $P > 0.05$ ) changes between them. The serum concentration of progesterone was however, not altered by methanol extract ( $318.001 \pm 2.26$  ng/

ml) relative to control ( $320.60 \pm 0.19$  ng/ml) rats producing only a 0.8% non-significant reduction but still caused a 65.63% and 65.34% significant ( $110.21 \pm 1.15$  ng/ml) decrease relative to control and methanol extract treated lactating rats ( $320.60 \pm 0.19$  ng/ml,  $318.00 \pm 2.26$  ng/ml) respectively. The results presented in **Figures 1 and 2** is the bar charts of hormone concentrations in the various treatments administered to experimental lactating and the non-lactating Albino rats. The chart collaborate the results in **Tables 2 and 3** already discussed in this study.

### Discussion

The result obtained did not establish a definite reduction pattern of these selected lactogenic hormones in non-lactating and lactating rats as methanolic crude extract produced significant reduction in serum concentrations of prolactin in both non-lactating and lactating rats. However, methanolic extract of *Aframomum melegueta* seed produced significant reduction in serum levels of prolactin and estradiol in both non-lactating and lactating rats. The lowered serum concentration of prolactin in rats treated with standard reference drug bromocriptine may not be unconnected to its action as dopamine agonist which inhibits the secretion of prolactin [14]. The drug did not only reduce the serum concentration of prolactin but went ahead to decrease that of estrogen and progesterone in both lactating and non-lactating rats relative to control treatment. Methanol extract of *Aframomum melegueta* seed produced a significant decrease in serum concentration of prolactin of wistar rats similar to bromocriptine. This probably may be due to presence of alkaloid in *Aframomum melegueta* seed as is the case with bromocriptine [19, 23]. The decreased serum concentration of prolactin observed in this study is in agreement with the report of [24, 25] who observed a significant reduction in serum level of prolactin of wistar non-lactating rats treated with Alligator pepper seed. The hypothesis of a regulatory role of placenta in pituitary prolactin and luteinizing hor-

## Methanolic extract of A.m on lactogenic hormones

**Table 2.** Serum concentrations of selected lactogenic hormones in non-lactating rats (NL)


Treatment groups			
Lactogenic hormones	Bromocriptine 0.11 mg/kg/day	Control 100 mg/kg/day	Methanol extract 100 mg/kg/day
Prolactin (ng/ml)	12.56±0.89 <sup>b</sup>	33.23±1.82 <sup>c</sup>	26.15±0.15 <sup>m</sup>
Estrogen (estradiol ng/ml)	2.81±0.12 <sup>c,m</sup>	3.05±0.06 <sup>c</sup>	2.68±0.08 <sup>m</sup>
Progesterone (ng/ml)	127.96±0.06 <sup>b</sup>	323.43±0.20 <sup>c</sup>	315.72±7.72 <sup>c</sup>

Means are ± SD. Mean values on the same row with different superscripts are significantly (P<0.05) different. b-Bromocriptine; c-Control; m-Methanol extract.

**Table 3.** Serum concentrations of selected lactogenic hormones in lactating (L) rats

Treatment groups			
Lactogenic hormones	Bromocriptine 0.11 mg/kg/day	Control 100 mg/kg/day	Methanol extract 100 mg/kg/day
Prolactin (ng/ml)	10.56±0.29 <sup>b</sup>	19.90±1.16 <sup>c</sup>	14.90±1.16 <sup>m</sup>
Estrogen (estradiol ng/ml)	2.41±0.80 <sup>b</sup>	3.17±0.09 <sup>c</sup>	2.40±0.03 <sup>b</sup>
Progesterone (ng/ml)	110.21±1.16 <sup>b</sup>	320.60±0.19 <sup>c</sup>	318.001±2.26 <sup>c</sup>

Means are ± SD. Mean values on the same row with different superscripts are significantly (P<0.05) different. b=bromocriptine; c=control treatment; m=methanol extract.




**Figure 1.** A-C. Hormonal response to various treatments administered to lactating Albino rats.

mone (LH) and release by hypothalamus-pituitary stimulation may have played out here [26]. Although prolactin under specific condition

such as pregnancy, lactation and estrus responds to particular physiological demand [27] but methanol seed extract of *A. melegueta*

## Methanolic extract of A.m on lactogenic hormones



**Figure 2.** A-C. Hormonal response to various treatments administered to non-lactating Albino rats.

caused significant ( $P < 0.05$ ) reduction of serum prolactin in both non-lactating and lactating rats mimicking the actions of bromocriptine which is the standard reference drug. Bromocriptine produced a generalized reduction in serum concentrations of the three hormones selected in this research in both non-lactating and lactating rats and this was supported by the report of [28] who reported a significant decrease of serum prolactin by bromocriptine in all estradiol treated rats while researching on the effect of bromocriptine on serum prolactin levels, pituitary weight and immune reaction of prolactin cells in estradiol treated ovariectomized rats. Methanol seed extract of *Aframomum melegueta* brought about a decrease in serum levels of estradiol in lactating rat. This finding may correlate with the report of [29] on stimulating mammary growth development by estradiol just like prolactin. Conversely the reduced levels of these hormones by the

extract manifest the anti-lactogenic effect of the extract. The antiestrogenic activities in this study with respect to normal control treatment is in consonance the report of [30] on screening for estrogenic and antiestrogenic activities of plants growing in Egypt and Thailand. The finding was collaborated by the work of [31] who reported that estrogen enhances prolactin secretion and that plasma concentration is normally higher in women than men. If estrogen enhances prolactin production and methanol seed extract of *A. melegueta* reduces the serum levels of estrogen and therefore disrupting the capacity to enhance prolactin production. This may probably contribute to low level of serum prolactin in this study. In other words, decreased concentration of estradiol may probably results in decreased plasma concentration of prolactin as is the case with methanol extract in this study. This is however, slightly different from the report of [32] who stated that

## Methanolic extract of A.m on lactogenic hormones

prolactin and estradiol synergize in producing mammary growth but estradiol antagonizes the milk producing effect of prolactin on the udder. Unlike bromocriptine methanol seed extract of *Aframomum melegueta* did not produce significant reduction in serum progesterone concentration of Albino rats used in this study for both non-lactating and lactating rats. This is at variance with the report of [33] on effect of saline extract of Alligator pepper on serum progesterone in pregnant Spray Dawley rats. The non-significant response of the extract to serum progesterone is advantageous bearing in mind that progesterone withdrawal is the trigger that initiates lactation [34]. These results did not differentiate much the lactating from non-lactating rats since the extracts produced significant reduction in the affected hormones of lactating as well as the non-lactating Wistar rats. No reason has been advanced for this. Generally, serum prolactin levels are higher during the early part of lactation and declines toward the end [35].

### Conclusion

Prolactin, estrogen and progesterone all play one role or the other in mammary gland development and thereby enhancing lactation. These roles are played by these hormones at different stages of mammary gland development. To create a ductal tree that fills the fat pad, branching morphogenesis is initiated at puberty by growth hormone and estrogen. Upon pregnancy, the action of progesterone and prolactin generate alveoli which secrete milk during lactation. Prolactin is the major generator of lactational competence during pregnancy and these functions both indirectly through its regulation of ovarian progesterone production and directly via its effects on mammary epithelia cells. Agonistic or antagonistic actions to these functions will either elevate or lower concentrations of these hormones thereby increasing or decreasing mammary gland development *viz*-*viz* milk secretion. The reduction in serum levels of prolactin and estrogen in both lactating and non-lactating Albino rats treated with methanolic seed extract of *Aframomum melegueta* demonstrated anti-lactogenesis even though serum progesterone was not significantly affected by the treatment. By the findings of this study, *Aframomum melegueta* seed extract slightly mimicked the reference drug bromocriptine and could be a good alternative in managing

cases of hyperprolactinemia and galactorrhoea in human.

### Acknowledgements

I acknowledge the work of Mr. Lucky Ebri who helped in typesetting the manuscript.

### Disclosure of conflict of interest

None.

**Address correspondence to:** Hyacinth Ibekwe, Department of Animal Science, Cross River University of Technology (CRUTECH) Obubra Campus, Nigeria. E-mail: hibekwe2010@gmail.com

### References

- [1] Wolfgang W, Hubertus J and Dana Seidlova W. Definition, classification and mechanism of action of endocrine disrupting chemicals. *Hormones Mini Review* 2010; 9: 15-20.
- [2] Neville MC, Morton J and Unemura S. Lactogenesis: transition from pregnancy to lactation. *Pediatr Clin North Am* 2001; 48: 35-52.
- [3] Falconer IR. Lactation: proceeding of international symposium, forming the 17th Easter School in Agricultural Science. Butterworths: University of Nottingham; 1970. pp. 173-175.
- [4] Cowie AT. Influence of hormones on mammary growth and milk secretion. In IR. Falconer's ed. *Lactation*. Nottingham: University of Nottingham Press; 1971. pp. 123-140.
- [5] Talwalker PK and Meites J. Tubuloalveolar developmental effects of selected anterior pituitary hormones. *Proceedings of the Society for Experimental Biology and Medicine* 1964; 117: 121-126.
- [6] Foltin J, Janene M, Jim R, Karen D. Endocrinology, metabolism and reproductive function. In WF. Ganong Ed. *Review of medical physiology*. 21st edition. New York: Lange Medical Books; 2003. pp. 282-490.
- [7] Melmed S and Kleinberg D. Anterior pituitary. In: Kroneberg HM, Melmed S, Polonsky KS and Larsen PR, editors. *Williams textbook of endocrinology*. Philadelphia: W. B. Saunders; 2008. pp. 200-218.
- [8] Fleet IR, Goode JA, Hamon MH, Laurice MS, Linzell JL, Peaker M. Secretory activity of goat mammary glands during pregnancy and lactation. *J Physiol* 1975; 251: 763-73.
- [9] Daryl KG. Hormones of the gonads. In: MKM Robert editor. *Harper's Biochemistry*. 24th edition. New York: Prentice Hall; 1996. pp. 566-580.
- [10] Stanley L. *Clinical laboratory tests, values and implications*. Pennsylvania: Springhouse Corporation; 2005. pp. 29-97.

## Methanolic extract of A.m on lactogenic hormones

- [11] Pagana KD and Pagana TJ. Mosby's manual of diagnostic and laboratory tests. 3rd edition. Mosby: St. Louis; 2006. pp. 114-118.
- [12] Clive P, Michael C, Morley S, Michael W and Brain H. Integrated pharmacology. 2nd edition. London: Mosby; 2002. pp. 281-329.
- [13] Lawrence RA and Lawrence RM. Breastfeeding: a guide for the medical profession. 5th edition. Mosby: St. Louis; 1999. pp. 118-128.
- [14] daSilva OP, Knoppert DC, Angelini MM and Forret P. Effect of domperidone on milk production in mothers of premature newborns: a randomized, double-blind placebo controlled trial. *CMAJ* 2001; 164: 17-21.
- [15] Paul EB. The dispensatory of the United State of America. The Grain of Paradise *Speculum* 1961; 36: 302-307.
- [16] Maguelonne TS and Anthea B Jr. Synonymous of melegueta pepper. *The History of Food* 2009; 1: 446-451.
- [17] John L. The herb book. New York: Bantam Books; 1984. pp. 180-197.
- [18] Henderson AJ and Faulkner A. The effects of vincristine on milk secretion and milk composition in the goat. *Q J Exp Physiol* 1985; 70: 15-22.
- [19] Okwu DE. Phytochemicals, vitamins and mineral contents of two Nigerian medicinal plants. *International Journal of Molecular Medicine and Advance Sciences* 2005; 1: 375-381.
- [20] Frantz AG. Enzyme immunosorbent assay for prolactin. *New England Journal of Medicine* 1978; 298: 201-205.
- [21] March CM, Geobelsmann U, Nakamura RM and Mishell DR Jr. Roles of estradiol and progesterone in eliciting the midcycle luteinizing hormone and follicle stimulating hormone surges. *J Clin Endocrinol Metab* 1979; 49: 507-13.
- [22] Radwanska E, Frankenberg J and Allen EI. Plasma progesterone levels in normal and abnormal early human pregnancy. *Fertil Steril* 1978; 30: 398-402.
- [23] Okoli RI, Aigbe O, Obodo O and Mensah JK. Medicinal herbs used in managing some common ailments among Esan people of Edo State, Nigeria. *Pakistan Journal of Nutrition* 2007; 6: 490-496.
- [24] Ebong PE, Eyong EU and Lawal BAS. Prolactin lowering effect of Alligator pepper *A. melegueta*). *West African Journal of Pharmacology* 2000; 8: 185-186.
- [25] Mark N. Cupboard love: a dictionary of culinary curiosities. Toronto: Insomniac Press; 2004. pp. 77-83.
- [26] Tripathi KD. Essentials of medical pharmacology. 4th edition. New Delhi: Jaypee Brothers Medical Publishers; 2001. pp. 239-332.
- [27] Bray JJ, Cragg PA, Macknight ADC, Mills RG and Taylor DW. Lecture notes on human physiology. London: Blackwell Scientific Publications; 1992. pp. 214-230.
- [28] Ribeiro MF, Spritzer PM, Barbosa-Coutinho LM, Oliveira MC, Pavanato MA, Silva SB and Reis FM. Effect of bromocriptine on serum prolactin levels, pituitary weight and immunoreactive prolactin cells in estradiol-treated ovariectomized rats: an experimental model of estrogen-dependent hyperprolactinemia. 1997.
- [29] Agarwal RK, Srivastava AK and Kumar K. *Animal Physiology and Biochemistry Revised Edition*. Ram Nagar, New Delhi: S.Chand and Company; 2014. pp. 177-211.
- [30] El-Halawany AM, El Dine RS, Chung MH, Nishihara T, Hattori M. Estrogenic and antiestrogenic activities of plants growing in Egypt and Thailand. *Pharmacognosy Res* 2011; 3: 107-13.
- [31] Hoehn K and Marieb EN. *Human Anatomy and Physiology*. San Francisco: Pearson Benjamin Cummings; 2007. pp. 605-606.
- [32] Ganong WF. *Review of medical physiology* 2003. New York: McGraw-Hill Medical Publishing Division; 2003. pp. 458-470.
- [33] Inegbenebor V and Ebomoyi M. Effect of saline extract of Alligator pepper (*Zingiberaceae Aframomum melegueta*) on serum progesterone in pregnant Sprague dawley rats. *International Journal of Herbs and pharmacology Research* 2014; 3: 46-51.
- [34] Bandyopadhyay SK, Bhattacharyya, Choudhury RR and Basu S. *Textbook of veterinary gynaecology, artificial insemination, obstetrics and assisted reproduction*. New Delhi: Kalyani Publishers; 2013. pp. 66-70.
- [35] Miller JW. Drugs and the endocrine and metabolic system. In: Clive P, Michael C, Morley S, Michael W and Brain H, editors. *Integrated pharmacology*. 2nd edition. London: Mosby; 2002. pp. 281-288.